

# Annals of Clinical and Medical Case Reports

Research Article

ISSN 2639-8109 | Volume 6

## Depression and Anxiety Frequency in Patients Hospitalized on the Guadalajara Regional Military Hospital in the Month of April 2019

Gabriel MN<sup>1\*</sup> and Gilberto GFL<sup>2</sup>

<sup>1</sup>Major of military armed forces of Mexico, Neurologist, Mexico

<sup>2</sup>Medical Doctor, Social Service, Lamar University, Guadalajara, Mexico

### \*Corresponding author:

Gabriel Miranda Nava,  
Neurologist Head of the Neurology Service of the  
Regional Military Specialty Hospital of  
Guadalajara, Jalisco, Mexico,  
E-mail: drgabrielmiranda@hotmail.com

Received: 27 Feb 2021

Accepted: 12 Mar 2021

Published: 18 Mar 2021

### Copyright:

©2021 Gabriel MN et al., This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

### Citation:

Gabriel MN. Depression and Anxiety Frequency in Patients Hospitalized on the Guadalajara Regional Military Hospital in the Month of April 2019. *Ann Clin Med Case Rep.* 2021; V6(4): 1-6

### 1. Abstract

Observe and Identify patients that presented depression and anxiety using the Hospital Anxiety Diagnosis Scale (HADS), Zung & Conde scale and ASQ 15 scale on Guadalajara Regional Military Hospital during the month of April 01st to April 30th, 2019. Methods: This is a cohort, non-experimental, observational, prospective, and longitudinal study with PubMed and NCBI articles as variables. Findings: Most patients presented anxiety, most patients had a chronic illness, depression was seen mostly in patients older than 50 years old, there was inadequate diet, lack of sleep, and low distress level.

### 2. Introduction

Depression and anxiety affect most people around the world, it is characterized by a presence of fear, loss of interest, feelings of guilt or self-esteem that are more commonly associated with sleep disorders, lack of appetite, lack of energy or difficulty concentrating. Depression can become chronic or recurrent and difficult the overall performance daily, or capacity to live day by day, in its most dangerous form it can lead often to suicide and its lowest form it can be treated with medication and professional psychotherapy, (salud, Depression, 2017).

Anxiety is one of the major disorders and its characterized by persistent concern during any activity or routine it is difficult to treat

and it can affect the way a person feels physically, (mayoclinic, 2018).


During this investigation we will observe a sample that was taken on the Guadalajara Regional Military Hospital during the month of April a sample of 56 patients presented anxiety and depression according to three scales that were applied [1].

#### 2.1. HADS (Hospital Anxiety Diagnosis Scale)

The Hospital Anxiety Diagnosis Scale is an auto applicable questionnaire integrated by 14 items with subscales of seven items one for impaired questions and one with pair questions for depression, the authors for this scale are Sigmund and Snaith who proposed this in 1983 and defined the concepts of anxiety and depression the objective of this scale is to identify if the patient has being tensed, concerned or frightened in any way, the 8 items that form the depression subscale are centered around anhedonia with a maximum score that binds from 0 to a 39 score, in which 0-9 score means lack of stress, 10-19 means low stress, 19 to 29 means mild stress and 30 to 39 means anxiety and severe depression [2].


#### 2.2. Zung & Conde Scale

Its and auto applicable scale consisting of 20 phrases related to depression formed by 10 negative phrases and 10 positive phrases which relate to strong somatic symptoms and 8 cognitive items for each group contemplating the scale with two items refer to mood and other psychotic symptoms, (Roza, 2019) (Figure 1-4).




**Figure 1:** Patients with depression

- 62% of the patients did not present depression 34 patients
- 16 patients had low depression 29%
- 8% of the patients had moderate depression which represented 4
- 2 of the patients representing 1% presented what could be considered as severe depression




**Figure 2:** Most common ages seen during study

- 42% of the patients presented depression were over 50 years old.
- 35% of the patients presented depression between 40-50 years of age.
- 17% of the patients that presented depression had between 30-40 years of age.
- 6% of the patients had between the ages of 20 to 30 years of age.



**Figure 3:** Patients that had some sort of stress evaluated by HADS scale

- 43% of the patients had lack of stress.
- 46% of the patients had low level stress.
- 11% of the patients had moderate stress.
- 0% of the patients had severe stress.



**Figure 4:** Most common diseases (12 patients did not have an illness)


- Hipertensión: 16
- Diabetes Mellitus:12
- Renal Insufficiency: 4
- Ulcerative Colitis:1
- Hepatic Cirrosis: 3
- Lung Cancer: 3
- Cervical Cancer:1
- Fractures: 4

**2.3. Depression and Anxiety**

Severe Depression: Its characterized by a combination of symptoms that interfere with capacity to work, sleep, study, eat and enjoy daily basis activities [3].


**2.4. Dysthimic Disorder**

Its characterized by symptoms that is somewhat between 2 years and beyond but less severe, it incapacitates the patient, and it prevents him from having a normal life accompanied by a severe depression episode during life (Figure 5,6).



**Figure 5:** Civil Status

- Married: 54%
- Separated: 35%
- Single: 11%



**Figure 6:** Patients that presented anxiety

- 64% presented anxiety 35 patients
- 36% did not present anxiety 21 patients

**2.5. Psychotic Depression**

Occurs during severe depression and its accompanied by some form of psychosis accompanied by delirium and hallucinations.


**2.6. Seasonal Depression**

Its characterized by depression that appears during Winter or times

of decreased sunlight.

**2.7. Bipolar Disorder**

Its characterized by maniac depression disorder that it accompanied by cyclic mood swings and depression state, its often seen in patients with cancer, HIV/Aids and Parkinson (Figure 7).



**Figure 7: Balanced Diet**

- 86% No
- 14% Yes

### 3. Symptoms

#### 3.1. Emotional

Are accompanied by guilt ideas, a severe disease, ideas of sadness never going to heal, loneliness, lack of concentration because patient will eventually die [4].

#### 3.2. Physical

Difficulty eating, or basic needs, weight loss, mood swings.

#### 3.3. Negative thoughts

This is mostly seen in older patients, self-esteem problems, most cases are seen in patients over 60 years old, or below 45 years old.

### 4. Methods

This is a cohort, non-experimental, observational, prospective, and longitudinal study in which scholarship was evaluated, cause of hospitalization, age, previous diseases, job, and the days patient had been hospitalized [5].

56 patients both men and women older than 18 years old were evaluated during this study, a random sample was taken in which every patient has the same possibility of presenting depression or anxiety [8]. Patients hospitalized in the Women's hospital room, the mens hospital room, and the room that consisted of patients that had the rank of major in the Mexican armed forces or above excluding patients that belonged to Intensive care unit, using the Hospital Anxiety Diagnosis Scale, Zung & Conde Scale and ASQ-15 Scale were used during this study [6].

### 5. Justification

This study was conducted to observe what was the impact of being hospitalized and the relation it had with depression and anxiety in patient, we pretended [9] to find viable date that allowed us to expose the hospital environment and the presence of disease, anxiety, and depression [10].

### 6. Results

Patients with depression, Fuente, [11] Zung & Conde Depression Scale. 62% of the patients did not present depression 34 patients, 16 patients had low depression 29%, also 8% of the patients had moderate depression which represented 4 patients also 2 of the patients representing 1% presented what could be considered as severe depression [7].

### 7. Conclusion

Referred to this subject we understand this two disorders are preventable and the patient if its treated and seeks help with time he can change his daily habits, our suggestions are that a stable lifestyle with a well-balanced diet consisting of fruit, vegetables, meat, daily exercise, stable relationships with family and friends, alongside no work stress, in addition to a good mental stability can lead to a good life and to prevent this type of disorders also to prevent chronic illness which were seen during this study on most patients that indicated feeling anxious or depressed, if they are in this state also to take medication on time and with the help of family members and friend [12].

- We observe most patients presented a low depression level
- Most patients presented anxiety
- Most patients had a chronic illness
- Patients over 50 years old presented higher depression levels
- Most patients were married
- Most patients had low level stress level.

### References:

1. Casanova C. Screening for symptoms of anxiety and depression in patients admitted to a university hospital with acute coronary syndrome. 2019.
2. Group of Trabajo de Guia de Práctica clínica for el manejo paciente con trastornos de ansiedad en atención primaria. Plan Nacional for SNS. 2008.
3. Servicio de Psiquiatria y Psicología Hospital Universitario Dexeus. Barcelona. 2016.
4. National Institute of Mental Health. Department of Salud and Servicios Humanos de los Estados Unidos. Trastornos de Ansiedad.
5. Enfocando la depresión como problema de salud pública en México Fernando A. Wagner. Catalina González. 2012.
6. Hernandez G, Gricel, Monica, Kimmelman, Orellana Rev. Fr. méd. Chile v.133 n.8 Santiago ago. 2005 Trastornos de ansiedad en pacientes hospitalizados en Medicina Interna. 2005.
7. Costas M, Prado V, Crespo JM. Anxious and depressed enter patient hospitalizations at the Ferrol Hospital. 2013.
8. Haas SA, Levandowski DC, Kalil AN. Anxious, depressed, perceived and sensitized with cancer indicators for cytogenetic file. 2017.
9. Ye Li, Mei-Rong, Lv Yan-Jin, Wei Ling, Zhang SJX, Zhang HG. Dietary patterns and depression risk. A meta-analysis. 2017.
10. General IMSS. Consejo de Salubridad Guia de referencia rápida guía práctica clínica.
11. Organización Mundial de la Salud. Who is depressed and answered.
12. Roza M. escala autoadministrada de depresión psicomag. Escala de Zung y Conde que es. 2019.